

Cut-free sequent calculi for logics with adjoint modalities

Mehrnoosh Sadrzadeh (Universities of Oxford & Paris VII)
and
Roy Dyckhoff (University of St Andrews)

ASL Logic Colloquium,
Bern, 3 July 2008

Outline of talk

1. Algebras (here, distributive lattices) with adjoint modalities
2. Gentzen and Belnap style sequent calculi
3. Rules of our calculus, soundness
4. Cut elimination argument
5. Consequences (completeness, decidability)
6. Conclusion and future plans

Algebras with adjoint modalities

Classical algebraic modal logic

Boolean Algebra with De Morgan dual operators (\Box, \Diamond)

Non-classical algebraic modal logic

weaken the base and the duality

Heyting algebra with 'dual' operators

Complete lattice with *adjoint* operators

Distributive Lattices with Adjoint Modalities (DLAM)

Let \mathcal{A} be a set, with elements called *agents*. A DLAM **over** \mathcal{A} is a bounded distributive lattice L with an \mathcal{A} -indexed family of maps $\{f_A\}_{A \in \mathcal{A}} : L \rightarrow L$, each with a right adjoint $\square_A : L \rightarrow L$. Thus, for all $l, l', (l_i : i \in I)$, the following hold (for finite I), :

$$f_A(l) \leq l' \quad \text{iff} \quad l \leq \square_A l'$$

$$f_A\left(\bigvee_i l_i\right) = \bigvee_i f_A(l_i)$$

$$\square_A\left(\bigwedge_i l_i\right) = \bigwedge_i \square_A(l_i)$$

In particular, $f_A \perp = \perp$ and $\square_A \top = \top$; and all f_A and \square_A are order-preserving. If the lattice is complete, then the existence of the right adjoints follows routinely provided the maps f_A (exist and) preserve arbitrary joins.

Application

Reasoning about knowledge of agents $A \in \mathcal{A}$

$m, m' \in M$: logical propositions

$m \vee m', m \wedge m'$: logical disjunction, conjunction

$m \leq m'$: logical consequence

$f_A(m)$: **appearance of agent A about m**

All the propositions that appear to agent A
to be true when m holds in reality.

$\square_A m$: agent A knows that m

Gentzen-style sequent calculi

Sequent calculi for reasoning about lattices are old; there are two forms. The first [4,5] considers sequents of “**atoms**” $t \leq t'$ and is useful to decide quasi-equations in lattice theory, e.g.

$$\frac{\Gamma \Rightarrow s \leq t \quad \Gamma \Rightarrow s \leq t'}{\Gamma \Rightarrow s \leq (t \wedge t')} R\wedge$$

The second kind (see [7]) considers sequents of **terms**, with the sequent arrow to represent the lattice order \leq , and, with

$$t = t' \text{ iff } t \Rightarrow t' \text{ and } t' \Rightarrow t,$$

decides the equational theory of **distributive** lattices, e.g. with

$$\frac{\Gamma \Rightarrow t \quad \Gamma \Rightarrow t'}{\Gamma \Rightarrow t \wedge t'} R\wedge$$

Ours is a generalisation of this second kind.

Gentzen-style sequent calculi, 2

[1] gives a Gentzen-style calculus (of the second kind) for modelling epistemic actions as resources. The calculus allows reasoning about “epistemic systems”, modelled by a quantale Q acting on a Q -module of epistemic propositions and facts.

The calculus includes a cut rule; this appears to be (in the context of that calculus) non-eliminable; we seek to remedy this as a basis for automation of proof search.

Belnap-style sequent calculi

Our calculus also includes agents, whose presence forces a richer structure of sequents. Moortgat [6] attributes this to Belnap [2], and exploits it (in a linguistic context) for residuated lattices with modal operators. (Our work is thus a variation on Moortgat's work, distinguished by organisation of rules to allow Weakening to be admissible.)

Syntax of our calculus: Formulae, items, contexts

The set M of *formulae* m of our language is generated over sets At of *atoms* p and \mathcal{A} of *agents* A and by the following grammar:

$$m ::= \perp \mid \top \mid p \mid m \wedge m \mid m \vee m \mid \Box_A m \mid f_A(m)$$

Items I, J, \dots and *contexts* Γ, Δ, \dots are generated by the following syntax:

$$\begin{aligned} I & ::= m \mid \Gamma^A \\ \Gamma & ::= \{I, I, \dots, I\} \end{aligned}$$

Thus, contexts are finite multisets of items, whereas items are either formulae or contexts annotated with an agent. (Where there is only one agent, the name *boxed context* can be used, as in [3].)

Syntax of our calculus: Contexts with holes

The notion of *context-with-a-hole* Δ is defined as follows:

$$\Delta ::= (\Gamma, []) \mid (\Gamma, \Delta^A)$$

and so a context-with-a-hole is a context (i.e. a multiset of items) together with either a hole or an agent-annotated context-with-a-hole. **Note that a context-with-a-hole is not a context.**

The result $\Delta[\Gamma]$ of *applying* Δ to a context Γ , replacing the hole $[]$ by Γ , is a context, defined recursively as follows:

$$\begin{aligned} (\Gamma, []) [\Gamma'] &= \Gamma, \Gamma' \\ (\Gamma, \Delta^A) [\Gamma'] &= \Gamma, \Delta[\Gamma']^A \end{aligned}$$

where the commas in the right-hand sides indicate multiset union. This will often be applied in the particular case where Γ' is a single item (treated as a one-element multiset).

Rules of our calculus: Axioms (initial sequents)

$$\boxed{\overline{\Gamma, m \vdash m} \text{ } Id \qquad \overline{\Delta[\perp] \vdash m} \perp L \qquad \overline{\Gamma \vdash \top} \top R}$$

The first rule is restricted to atoms m ; but it can be shown that all sequents $\Gamma, m \vdash m$ are derivable. Note that the $\perp L$ rule allows the \perp to appear anywhere (as an item) deep inside the context, whereas the Id rule requires the principal formula to appear at top level in the context. This captures the requirement that (in the lattice interpretation) $f_A(\perp) = \perp$ (this is not generally true for arbitrary elements).

Rules of our calculus: Rules for the lattice operations

$$\frac{\Delta[m_1, m_2] \vdash m}{\Delta[m_1 \wedge m_2] \vdash m} \wedge L$$

$$\frac{\Gamma \vdash m_1 \quad \Gamma \vdash m_2}{\Gamma \vdash m_1 \wedge m_2} \wedge R$$

$$\frac{\Delta[m_1] \vdash m \quad \Delta[m_2] \vdash m}{\Delta[m_1 \vee m_2] \vdash m} \vee L$$

$$\frac{\Gamma \vdash m_i}{\Gamma \vdash m_1 \vee m_2} \vee R_i$$

Rules of our calculus: Rules for the modalities

$$\frac{\Delta[m^A] \vdash m'}{\Delta[f_A(m)] \vdash m'} f_{AL} \qquad \frac{\Gamma \vdash m}{\Gamma', \Gamma^A \vdash f_A(m)} f_{AR}$$

$$\frac{\Delta[(\Box_A m)^A, m] \vdash m'}{\Delta[(\Box_A m)^A] \vdash m'} \Box_{AL} \qquad \frac{\Gamma^A \vdash m}{\Gamma \vdash \Box_A m} \Box_{AR}$$

Rules of our calculus:

Interaction between exponentiation and meet

Finally, we need a rule (named K , following [6])

$$\boxed{\frac{\Delta[(\Gamma', \Gamma'')^A, \Gamma'^A, \Gamma''^A] \vdash m}{\Delta[(\Gamma', \Gamma'')^A] \vdash m} K}$$

Example Derivations

$$\frac{\frac{\overline{m \vdash m}}{m^A \vdash f_A(m)} f_A R}{m \vdash \Box_A f_A(m)} \Box_A R$$

$$\frac{\overline{(\Box_A m)^A, m \vdash m}}{(\Box_A m)^A \vdash m} \Box_A L \quad \frac{(\Box_A m)^A \vdash m}{f_A(\Box_A m) \vdash m} f_A L$$

$$\frac{\frac{\frac{\overline{m \vdash m}}{m^A \vdash f_A(m)} f_A R}{m^A \vdash f_A(m) \vee f_A(m')} \vee R \quad \frac{\frac{\overline{m' \vdash m'}}{m'^A \vdash f_A(m')} f_A R}{m'^A \vdash f_A(m) \vee f_A(m')} \vee R}{\frac{(m \vee m')^A \vdash f_A(m) \vee f_A(m')}{f_A(m \vee m') \vdash f_A(m) \vee f_A(m')} f_A L} \vee R$$

(Converse of the last is easy.)

Example Derivations, 2

$$\begin{array}{c}
 \frac{\overline{m \vdash m}}{\dots, m^A, m'^A \vdash f_A(m)} f_A R \quad \frac{\overline{m' \vdash m'}}{\dots, m^A, m'^A \vdash f_A(m')} f_A R \\
 \hline
 \dots, m^A, m'^A \vdash f_A(m) \wedge f_A(m') \quad \wedge R \\
 \hline
 \dots, (m \wedge m')^A \vdash f_A(m) \wedge f_A(m') \quad K \\
 \hline
 f_A(m \wedge m') \vdash f_A(m) \wedge f_A(m') \quad f_A L
 \end{array}$$

Semantics of sequents; soundness:

Let L be a DLAM over \mathcal{A} . An **interpretation** of the set M of formulae (over the atoms At and agents \mathcal{A}) in L is a map: $\llbracket - \rrbracket: At \rightarrow L$.

Meaning of a formula: by induction on the structure, e.g.

$$\llbracket m_1 \wedge m_2 \rrbracket = \llbracket m_1 \rrbracket \wedge \llbracket m_2 \rrbracket, \quad \llbracket f_A(m) \rrbracket = f_A(\llbracket m \rrbracket)$$

Meaning of an item

$$\llbracket m \rrbracket = \text{as above}, \quad \llbracket \Gamma^A \rrbracket = f_A(\llbracket \Gamma \rrbracket)$$

Meaning of a context

$$\llbracket \{I_1, \dots, I_n\} \rrbracket = \llbracket I_1 \rrbracket \wedge \dots \wedge \llbracket I_n \rrbracket$$

Semantics of sequents; soundness:

Let L be a DLAM.

Definition. *Truth.*

A sequent $\Gamma \vdash m$ is *true in an interpretation* $\llbracket - \rrbracket$ in L iff $\llbracket \Gamma \rrbracket \leq \llbracket m \rrbracket$.

A sequent $\Gamma \vdash m$ is *true in L* iff true in every interpretation in L .

Definition. *Satisfiability.*

A sequent $\Gamma \vdash m$ is *satisfiable* (in L) iff there is an interpretation (in L) in which it is true.

Definition. *Validity.*

A sequent $\Gamma \vdash m$ is *valid* iff it is true in all interpretations.

Theorem. *Soundness.*

Any derivable sequent is valid.

Admissible Structural Rules

Lemma. The following *Weakening* rule is dp-admissible

$$\frac{\Delta[\Gamma] \vdash m}{\Delta[\Gamma, \Gamma'] \vdash m} Wk$$

Proof. Induction on the height of the derivation of the premiss and case analysis. E.g., suppose the last step is by $f_A R$, with $m = f_A m'$, and with premiss $\Gamma^* \vdash m'$ and conclusion $\Gamma'', \Gamma^{*A} \vdash m$, so $\Delta[\Gamma] = \Gamma'', \Gamma^{*A}$. To obtain $\Delta[\Gamma, \Gamma']$ from this there are two possibilities. In the first case, Γ occurs inside Γ^{*A} , and we make a routine use of the inductive hypothesis and reapply $f_A R$ with the same parameter. In the second case, we just use the $f_A R$ rule with a different parameter.

Admissible Structural Rules, 2

The dp-admissibility is used when we consider the K rule, since the place where the extra context Γ' is included may, in the premiss of that rule, be duplicated, requiring two weakenings by the inductive hypothesis.

Since (in the definition of 'context') we are using multi-sets rather than sets, there is a need to show admissibility of Contraction. This can be done; note the various duplications in the rules to ensure this works. (The rule is that from $\Delta[\Gamma, \Gamma] \vdash m$ we can infer $\Delta[\Gamma] \vdash m$.)

Cut elimination

Theorem. The *Cut* rule is admissible

$$\frac{\Gamma \vdash m \quad \Delta[m] \vdash m'}{\Delta[\Gamma] \vdash m'} \text{Cut}$$

Proof. Strong induction on the rank of the cut, where the *rank* is given by the pair (size of cut formula m , sum of heights of derivations of premisses).

Cut elimination: example step

Suppose the cut formula is $f_A(m)$ and is principal in both premisses:

$$\frac{\frac{\Gamma \vdash m}{\Gamma', \Gamma^A \vdash f_A(m)} f_{AR} \quad \frac{\Delta[m^A] \vdash m'}{\Delta[f_A(m)] \vdash m'} f_{AL}}{\Delta[\Gamma', \Gamma^A] \vdash m'} \text{Cut}$$

transforms to

$$\frac{\Gamma \vdash m \quad \Delta[m^A] \vdash m'}{\Delta[\Gamma^A] \vdash m'} \text{Cut}}{\Delta[\Gamma', \Gamma^A] \vdash m'} Wk$$

Consequences (transitivity)

Theorem. From $m \vdash m'$ and $m' \vdash m''$ follows $m \vdash m''$.

Consequences (completeness)

(Fix a set of atoms.)

Theorem. Let m, m' be formulae (in a language over the set \mathcal{A} of agents). The following are equivalent:

1. $m \vdash m'$;
2. $m \leq m'$ is true in all DLAMs over \mathcal{A} ;
3. $m \leq m'$ is true in all complete DLAMs over \mathcal{A} .

Thus, the sequent calculus is (w.r.t. the given semantics) sound and complete.

Consequences (proof of completeness, 3 implies 1)

Routine Lindenbaum-Tarski construction and completion.

Consequences (decidability)

Straightforward, using the sub-formula property and a loop-checker.
Is there a simple variant that allows avoidance of a loop-checker?

Conclusion and future plans

1. changing the representation, e.g. lists rather than multisets
2. proof-theoretic and implementation issues, e.g. invertibility lemmas, termination, loop-checking
3. enriching the base: Heyting algebras, quantales (for Linear Logic), systems [1]
4. comparing with other calculi for modal logics, e.g. deep inference systems (Guglielmi, Stouppa, Stewart, Brünnler, . . .)

References

1. Baltag, Coecke, Sadrzadeh. Epistemic actions as resources, *J. of Logic and Computation* 2007.
2. Belnap. Display Logic, *Journal of Phil. Logic* 1982.
3. K. Brünnler. Deep Sequent Systems for Modal Logic, *Advances in Modal Logic*, 2006.
4. RD, Negri. Decision methods for linearly ordered Heyting algebras, *Arch. Math. Logic* 2005.
5. Jipsen. Lattices, <http://math.chapman.edu/cgi-bin/structures/>.
6. Moortgat. Multimodal linguistic inference, *Logic J. of IGPL* 1995.
7. Ono, Komori. Logics without the contraction rule, *JSL* 1985
8. Troelstra, Schwichtenberg. Basic Proof Theory, CUP 2001.