

# $\omega$ -Degree Spectra

Alexandra A. Soskova

Sofia University

LC 2008

Bern 03.07.08

# Outline

- ▶ Degree spectra and jump spectra
- ▶  $\omega$ -enumeration degrees
- ▶  $\omega$ -degree spectra
- ▶  $\omega$ -co-spectra
- ▶ A minimal pair theorem
- ▶ Quasi-minimal degrees

# Enumeration of a Structure

Let  $\mathfrak{A} = (\mathbb{N}; R_1, \dots, R_k, =, \neq)$  be a countable abstract structure.

- ▶ An enumeration  $f$  of  $\mathfrak{A}$  is a total mapping from  $\mathbb{N}$  onto  $\mathbb{N}$ .
- ▶ for any  $A \subseteq \mathbb{N}^a$  let
$$f^{-1}(A) = \{\langle x_1 \dots x_a \rangle : (f(x_1), \dots, f(x_a)) \in A\}.$$
- ▶  $f^{-1}(\mathfrak{A}) = f^{-1}(R_1) \oplus \dots \oplus f^{-1}(R_k) \oplus f^{-1}(=) \oplus f^{-1}(\neq).$

## Definition (L. Richter, 1981)

The Turing degree spectrum of  $\mathfrak{A}$

$$DS_T(\mathfrak{A}) = \{d_T(f^{-1}(\mathfrak{A})) \mid f \text{ is an injective enumeration of } \mathfrak{A}\}$$

- ▶ J. Knight, Ash, Jockush, Downey, Slaman.

# Degree Spectra and Co-spectra

$\omega$ -Degree Spectra

Alexandra A.  
Soskova

Degree Spectra

$\omega$ -Enumeration  
Degrees

$\omega$ -Degree Spectra

Properties of the  
 $\omega$ -Degree Spectra

Minimal Pair Theorem  
Quasi-Minimal Degree

## Definition (Soskov, 2004)

- ▶ The degree spectrum of  $\mathfrak{A}$

$$DS(\mathfrak{A}) = \{d_e(f^{-1}(\mathfrak{A})) \mid f \text{ is an enumeration of } \mathfrak{A}\}.$$

- ▶ The co-spectrum of  $\mathfrak{A}$

$$CS(\mathfrak{A}) = \{\mathbf{b} : (\forall \mathbf{a} \in DS(\mathfrak{A}))(\mathbf{b} \leq \mathbf{a})\}.$$

# Degree Spectra and Co-spectra

$\omega$ -Degree Spectra

Alexandra A.  
Soskova

Degree Spectra

$\omega$ -Enumeration  
Degrees

$\omega$ -Degree Spectra

Properties of the  
 $\omega$ -Degree Spectra

Minimal Pair Theorem  
Quasi-Minimal Degree

## Definition

Let  $\mathcal{A} \subseteq \mathcal{D}_e$ .  $\mathcal{A}$  is *upwards closed with respect to total enumeration degrees*, if

$$\mathbf{a} \in \mathcal{A}, \mathbf{b} \text{ is total and } \mathbf{a} \leq \mathbf{b} \Rightarrow \mathbf{b} \in \mathcal{A}.$$

The degree spectra are upwards closed with respect to total enumeration degrees.

# Properties of upwards closed sets

Let  $\mathcal{A} \subseteq \mathcal{D}_e$  be upwards closed with respect to total enumeration degrees. Denote by

$$\text{co}(\mathcal{A}) = \{b : b \in \mathcal{D}_e \text{ \& } (\forall a \in \mathcal{A})(b \leq_e a)\}.$$

- ▶  $\mathcal{A}_t = \{\mathbf{a} : \mathbf{a} \in \mathcal{A} \text{ \& } \mathbf{a} \text{ is total}\} \implies \text{co}(\mathcal{A}) = \text{co}(\mathcal{A}_t).$
- ▶ Let  $\mathbf{b} \in \mathcal{D}_e$  and  $n > 0$ .

$$\mathcal{A}_{\mathbf{b},n} = \{\mathbf{a} : \mathbf{a} \in \mathcal{A} \text{ \& } \mathbf{b} \leq \mathbf{a}^{(n)}\} \implies \text{co}(\mathcal{A}) = \text{co}(\mathcal{A}_{\mathbf{b},n}).$$

# Properties of degree spectra and co-spectra

- ▶ Let  $\mathbf{c} \in DS_n(\mathfrak{A})$  and  $n > 0$ . Then

$$CS(\mathfrak{A}) = co(\{\mathbf{a} \mid \mathbf{a} \in DS(\mathfrak{A}) \ \& \ \mathbf{a}^{(n)} = \mathbf{c}\}).$$

- ▶ A minimal pair theorem:  
There exist  $\mathbf{f}$  and  $\mathbf{g}$  in  $DS(\mathfrak{A})$ :

$$(\forall \mathbf{a} \in \mathcal{D}_e)(\forall k)(\mathbf{a} \leq_e \mathbf{f}^{(k)} \ \& \ \mathbf{a} \leq_e \mathbf{g}^{(k)} \Rightarrow \mathbf{a} \in CS_k(\mathfrak{A})).$$

- ▶ Quasi-minimal degree:  
There exists  $\mathbf{q}_0$  quasi-minimal for  $DS(\mathfrak{A})$ 
  - ▶  $\mathbf{q}_0 \notin CS(\mathfrak{A})$ ;
  - ▶ for every total  $e$ -degree  $\mathbf{a}$ :  $\mathbf{a} \geq_e \mathbf{q}_0 \Rightarrow \mathbf{a} \in DS(\mathfrak{A})$  and  $\mathbf{a} \leq_e \mathbf{q}_0 \Rightarrow \mathbf{a} \in CS(\mathfrak{A})$ .
- ▶ Every countable ideal can be represented as a co-spectrum of some structure  $\mathfrak{A}$ .



# $\omega$ -Enumeration Degrees

- ▶ Uniform reducibility on sequences of sets
- ▶  $S$  the set of all sequences of sets of natural numbers
- ▶ For  $\mathcal{B} = \{B_n\}_{n < \omega} \in S$  call *the jump class of  $\mathcal{B}$*  the set

$$J_{\mathcal{B}} = \{d_T(X) \mid (\forall n)(B_n \text{ is c.e. in } X^{(n)} \text{ uniformly in } n)\} .$$

- ▶  $\mathcal{A} \leq_{\omega} \mathcal{B}$  ( $\mathcal{A}$  is  $\omega$ -enumeration reducible to  $\mathcal{B}$ ) if  $J_{\mathcal{B}} \subseteq J_{\mathcal{A}}$
- ▶  $\mathcal{A} \equiv_{\omega} \mathcal{B}$  if  $J_{\mathcal{A}} = J_{\mathcal{B}}$ .

- ▶  $\equiv_\omega$  is an equivalence relation on  $\mathcal{S}$ .
- ▶  $d_\omega(\mathcal{B}) = \{\mathcal{A} \mid \mathcal{A} \equiv_\omega \mathcal{B}\}$
- ▶  $\mathcal{D}_\omega = \{d_\omega(\mathcal{B}) \mid \mathcal{B} \in \mathcal{S}\}$ .
- ▶ If  $A \subseteq \mathbb{N}$  denote by  $A \uparrow \omega = \{A, \emptyset, \emptyset, \dots\}$ .
- ▶ For every  $A, B \subseteq \mathbb{N}$ :

$$A \leq_e B \iff A \uparrow \omega \leq_\omega B \uparrow \omega.$$

- ▶ The mapping  $\kappa(d_e(A)) = d_\omega(A \uparrow \omega)$  gives an isomorphic embedding of  $\mathcal{D}_e$  to  $\mathcal{D}_\omega$ .

# $\omega$ -Enumeration Degrees

$\omega$ -Degree Spectra

Alexandra A.  
Soskova

Degree Spectra

$\omega$ -Enumeration  
Degrees

$\omega$ -Degree Spectra

Properties of the  
 $\omega$ -Degree Spectra

Minimal Pair Theorem  
Quasi-Minimal Degree

Let  $\mathcal{B} = \{B_n\}_{n < \omega} \in \mathcal{S}$ .

A jump sequence  $\mathcal{P}(\mathcal{B}) = \{\mathcal{P}_n(\mathcal{B})\}_{n < \omega}$ :

1  $\mathcal{P}_0(\mathcal{B}) = B_0$

2  $\mathcal{P}_{n+1}(\mathcal{B}) = (\mathcal{P}_n(\mathcal{B}))' \oplus B_{n+1}$

**Theorem (Soskov, Kovachev)**

$A \leq_{\omega} B$ , if  $A_n \leq_e \mathcal{P}_n(\mathcal{B})$  uniformly in  $n$ .

# $\omega$ -Enumeration Jump

- ▶ For every  $\mathcal{A} \in \mathcal{S}$  the  $\omega$ -enumeration jump of  $\mathcal{A}$  is  $\mathcal{A}' = \{\mathcal{P}_{n+1}(\mathcal{A})\}_{n < \omega}$
- ▶  $d_\omega(\mathcal{A})' = d_\omega(\mathcal{A}')$
- ▶  $\mathcal{A}^{(k+1)} = (\mathcal{A}^{(k)})'$
- ▶  $d_\omega(\mathcal{A})^{(k+1)} = d_\omega(\mathcal{A}^{(k+1)})$
- ▶  $\mathcal{A}^{(k)} = \{\mathcal{P}_{n+k}(\mathcal{A})\}_{n < \omega}$  for each  $k$ .

Let  $\mathfrak{A}_1, \dots, \mathfrak{A}_n$  be given structures.

## Definition

The *relative spectrum*  $RS(\mathfrak{A}, \mathfrak{A}_1, \dots, \mathfrak{A}_n)$  of the structure  $\mathfrak{A}$  with respect to  $\mathfrak{A}_1, \dots, \mathfrak{A}_n$  is the set

$$\{d_e(f^{-1}(\mathfrak{A})) \mid f \text{ is an enumeration of } \mathfrak{A} \text{ \& } (\forall k \leq n)(f^{-1}(\mathfrak{A}_k) \leq_e f^{-1}(\mathfrak{A})^{(k)})\}$$

It turns out that all properties of the degree spectra remain true for the relative spectra.

Let  $\mathcal{B} = \{B_n\}_{n < \omega}$  be a fixed sequence of sets.

The enumeration  $f$  of the structure  $\mathfrak{A}$  is *acceptable with respect to  $\mathcal{B}$* , if for every  $n$ ,

$$f^{-1}(B_n) \leq_e f^{-1}(\mathfrak{A})^{(n)} \text{ uniformly in } n.$$

Denote by  $\mathcal{E}(\mathfrak{A}, \mathcal{B})$  - the class of all acceptable enumerations.

## Definition

The  $\omega$ -degree spectrum of  $\mathfrak{A}$  with respect to  $\mathcal{B} = \{B_n\}_{n < \omega}$  is the set

$$\text{DS}(\mathfrak{A}, \mathcal{B}) = \{d_e(f^{-1}(\mathfrak{A})) \mid f \in \mathcal{E}(\mathfrak{A}, \mathcal{B})\}$$

- ▶ It is easy to find a structure  $\mathfrak{A}$  and a sequence  $\mathcal{B}$  such that  $DS(\mathfrak{A}, \mathcal{B}) \neq DS(\mathfrak{A})$ .
- ▶ The notion of the  $\omega$ -degree spectrum is a generalization of the relative spectrum:  
 $RS(\mathfrak{A}, \mathfrak{A}_1, \dots, \mathfrak{A}_n) = DS(\mathfrak{A}, \mathcal{B})$ , where  $\mathcal{B} = \{B_k\}_{k < \omega}$ ,
  - ▶  $B_0 = \emptyset$ ,
  - ▶  $B_k$  is the positive diagram of the structure  $\mathfrak{A}_k$ ,  $k \leq n$
  - ▶  $B_k = \emptyset$  for all  $k > n$ .

# $\omega$ -Degree Spectra and Jump Spectra

## Proposition

$DS(\mathfrak{A}, \mathcal{B})$  is upwards closed with respect to total  $e$ -degrees.

## Definition

The  $k$ th  $\omega$ -jump spectrum of  $\mathfrak{A}$  with respect to  $\mathcal{B}$  is the set

$$DS_k(\mathfrak{A}, \mathcal{B}) = \{\mathbf{a}^{(k)} \mid \mathbf{a} \in DS(\mathfrak{A}, \mathcal{B})\}.$$

## Proposition

$DS_k(\mathfrak{A}, \mathcal{B})$  is upwards closed with respect to total  $e$ -degrees.



For every  $\mathcal{A} \subseteq \mathcal{D}_\omega$  let

$$\text{co}(\mathcal{A}) = \{\mathbf{b} \mid \mathbf{b} \in \mathcal{D}_\omega \ \& \ (\forall \mathbf{a} \in \mathcal{A})(\mathbf{b} \leq_\omega \mathbf{a})\}.$$

## Definition

The  $\omega$ -co-spectrum of  $\mathfrak{A}$  with respect to  $\mathcal{B}$  is the set

$$\text{CS}(\mathfrak{A}, \mathcal{B}) = \text{co}(\text{DS}(\mathfrak{A}, \mathcal{B})).$$

## Definition

The  $k$ th  $\omega$ -co-spectrum of  $\mathfrak{A}$  with respect to  $\mathcal{B}$  is the set

$$\text{CS}_k(\mathfrak{A}, \mathcal{B}) = \text{co}(\text{DS}_k(\mathfrak{A}, \mathcal{B})).$$

# Normal Form Theorem

Let  $\mathcal{L}$  be the language of the structure  $\mathfrak{A}$ . For each  $n$  let  $P_n$  be a new unary predicate representing the set  $B_n$ .

- ▶ An elementary  $\Sigma_0^+$  formula is an existential formula of the form  $\exists Y_1 \dots \exists Y_m \Phi(W_1, \dots, W_r, Y_1, \dots, Y_m)$ , where  $\Phi$  is a finite conjunction of atomic formulae in  $\mathcal{L} \cup \{P_0\}$ ;
- ▶ A  $\Sigma_n^+$  formula is a c.e. disjunction of elementary  $\Sigma_n^+$  formulae;
- ▶ An elementary  $\Sigma_{n+1}^+$  formula is a formula of the form  $\exists Y_1 \dots \exists Y_m \Phi(W_1, \dots, W_r, Y_1, \dots, Y_m)$ , where  $\Phi$  is a finite conjunction of atoms of the form  $P_{n+1}(Y_j)$  or  $P_{n+1}(W_i)$  and  $\Sigma_n^+$  formulae or negations of  $\Sigma_n^+$  formulae in  $\mathcal{L} \cup \{P_0\} \cup \dots \cup \{P_n\}$ .

# Normal Form Theorem

## Definition

The sequence  $\mathcal{A} = \{A_n\}_{n < \omega}$  is *formally  $k$ -definable* on  $\mathfrak{A}$  with respect to  $\mathcal{B}$  if there exists a computable sequence  $\{\Phi^{\gamma(n,x)}(W_1, \dots, W_r)\}_{n,x < \omega}$  of  $\Sigma_{n+k}^+$  formulae and elements  $t_1, \dots, t_r$  of  $\mathbb{N}$  such that for every  $x \in \mathbb{N}$ , the following equivalence holds:

$$x \in A_n \iff (\mathfrak{A}, \mathcal{B}) \models \Phi^{\gamma(n,x)}(W_1/t_1, \dots, W_r/t_r).$$

## Theorem

*The sequence  $\mathcal{A}$  is formally  $k$ -definable on  $\mathfrak{A}$  with respect to  $\mathcal{B}$  iff  $d_\omega(\mathcal{A}) \in \text{CS}_k(\mathfrak{A}, \mathcal{B})$ .*

# Properties of upwards closed sets

Let  $\mathcal{A} \subseteq \mathcal{D}_e$  be an upwards closed set with respect to total e-degrees.

## Proposition

$co(\mathcal{A}) = co(\{\mathbf{a} : \mathbf{a} \in \mathcal{A} \ \& \ \mathbf{a} \text{ is total}\})$ .

## Corrolary

$CS(\mathcal{A}, \mathcal{B}) = co(\{\mathbf{a} \mid \mathbf{a} \in DS(\mathcal{A}, \mathcal{B}) \ \& \ \mathbf{a} \text{ is a total e-degree}\})$ .

# Negative results (Vatev)

Let  $\mathcal{A} \subseteq \mathcal{D}_e$  be an upwards closed set with respect to total e-degrees and  $k > 0$ .

- ▶ There exists  $\mathbf{b} \in \mathcal{D}_e$  such that

$$co(\mathcal{A}) \neq co(\{\mathbf{a} : \mathbf{a} \in \mathcal{A} \ \& \ \mathbf{b} \leq \mathbf{a}^{(k)}\}).$$

- ▶ Let  $n > 0$ . There is a structure  $\mathfrak{A}$ , a sequence  $\mathcal{B}$  and  $\mathbf{c} \in DS_n(\mathfrak{A}, \mathcal{B})$  such that

$$CS(\mathfrak{A}, \mathcal{B}) \neq co(\{\mathbf{a} \in DS(\mathfrak{A}, \mathcal{B}) \mid \mathbf{a}^{(n)} = \mathbf{c}\}).$$

## Theorem

*For every structure  $\mathfrak{A}$  and every sequence  $\mathcal{B} \in \mathcal{S}$  there exist total enumeration degrees  $\mathbf{f}$  and  $\mathbf{g}$  in  $\text{DS}(\mathfrak{A}, \mathcal{B})$  such that for every  $\omega$ -enumeration degree  $\mathbf{a}$  and  $k \in \mathbb{N}$ :*

$$\mathbf{a} \leq_{\omega} \mathbf{f}^{(k)} \ \& \ \mathbf{a} \leq_{\omega} \mathbf{g}^{(k)} \Rightarrow \mathbf{a} \in \text{CS}_k(\mathfrak{A}, \mathcal{B}) .$$

## Corrolary

$CS_k(\mathfrak{A}, \mathfrak{B})$  is the least ideal containing all  $k$ th  $\omega$ -jumps of the elements of  $CS(\mathfrak{A}, \mathfrak{B})$ .

- ▶  $I = CS(\mathfrak{A}, \mathfrak{B})$  is a countable ideal;
- ▶  $CS(\mathfrak{A}, \mathfrak{B}) = I(\mathbf{f}) \cap I(\mathbf{g})$ ;
- ▶  $I^{(k)}$  - the least ideal, containing all  $k$ th  $\omega$ -jumps of the elements of  $I$ ;
- ▶ (Ganchev)  
 $I = I(\mathbf{f}) \cap I(\mathbf{g}) \implies I^{(k)} = I(\mathbf{f}^{(k)}) \cap I(\mathbf{g}^{(k)})$  for every  $k$ ;
- ▶  $I(\mathbf{f}^{(k)}) \cap I(\mathbf{g}^{(k)}) = CS_k(\mathfrak{A}, \mathfrak{B})$  for each  $k$
- ▶ Thus  $I^{(k)} = CS_k(\mathfrak{A}, \mathfrak{B})$ .

# Countable ideals of $\omega$ -enumeration degrees

There is a countable ideal  $I$  of  $\omega$ -enumeration degrees for which there is no structure  $\mathfrak{A}$  and sequence  $\mathcal{B}$  such that  $I = \text{CS}(\mathfrak{A}, \mathcal{B})$ .

- ▶  $\mathcal{A} = \{\mathbf{0}, \mathbf{0}', \mathbf{0}'', \dots, \mathbf{0}^{(n)}, \dots\}$ ;
- ▶  $I = I(\mathcal{A}) = \{\mathbf{a} \mid \mathbf{a} \in \mathcal{D}_\omega \ \& \ (\exists n)(\mathbf{a} \leq_\omega \mathbf{0}^{(n)})\}$  - a countable ideal generated by  $\mathcal{A}$ .
- ▶ Assume that there is a structure  $\mathfrak{A}$  and a sequence  $\mathcal{B}$  such that  $I = \text{CS}(\mathfrak{A}, \mathcal{B})$
- ▶ Then there is a minimal pair  $\mathbf{f}$  and  $\mathbf{g}$  for  $\text{DS}(\mathfrak{A}, \mathcal{B})$ , so  $I^{(n)} = I(\mathbf{f}^{(n)}) \cap I(\mathbf{g}^{(n)})$  for each  $n$ .
- ▶  $\mathbf{f} \geq \mathbf{0}^{(n)}$  and  $\mathbf{g} \geq \mathbf{0}^{(n)}$  for each  $n$ .
- ▶ Then by Enderton and Putnam [1970], Sacks [1971]:  $\mathbf{f}'' \geq \mathbf{0}^{(\omega)}$  and  $\mathbf{g}'' \geq \mathbf{0}^{(\omega)}$ .
- ▶ Hence  $I'' \neq I(\mathbf{f}'') \cap I(\mathbf{g}'')$ . A contradiction.









## Theorem

*For every structure  $\mathfrak{A}$  and every sequence  $\mathcal{B}$ , there exists  $F \subseteq \mathbb{N}$ , such that  $\mathbf{q} = d_\omega(F \uparrow \omega)$  and:*

- 1.  $\mathbf{q} \notin \text{CS}(\mathfrak{A}, \mathcal{B})$ ;*
- 2. If  $\mathbf{a}$  is a total e-degree and  $\mathbf{a} \geq_\omega \mathbf{q}$  then  $\mathbf{a} \in \text{DS}(\mathfrak{A}, \mathcal{B})$*
- 3. If  $\mathbf{a}$  is a total e-degree and  $\mathbf{a} \leq_\omega \mathbf{q}$  then  $\mathbf{a} \in \text{CS}(\mathfrak{A}, \mathcal{B})$ .*

## ► Questions:

- Is it true that for every structure  $\mathfrak{A}$  and every sequence  $\mathcal{B}$  there exists a structure  $\mathfrak{B}$  such that  $DS(\mathfrak{B}) = DS(\mathfrak{A}, \mathcal{B})$ ?
- If for a countable ideal  $I \subseteq \mathcal{D}_\omega$  there is an exact pair then are there a structure  $\mathfrak{A}$  and a sequence  $\mathcal{B}$  so that  $CS(\mathfrak{A}, \mathcal{B}) = I$ ?

-  Ganchev, H., Exact pair theorem for the  $\omega$ -enumeration degrees, *LNCS*, (B. Löwe S. B. Cooper and A. Sorbi, eds.), 4497, 316–324 (2007)
-  Soskov I. N., Degree spectra and co-spectra of structures. *Ann. Univ. Sofia*, 96 45–68, (2003).
-  Soskov, I. N., Kovachev, B. Uniform regular enumerations *Mathematical Structures in Comp. Sci.* 16 no. 5, 901–924, (2006)
-  Soskov, I. N. The  $\omega$ -enumeration degrees, *J. Logic and Computation* 17 no. 6, 1193-1214 (2007)
-  Soskov, I. N., Ganchev H. The jump operator on the  $\omega$ -enumeration degrees. *Ann. Pure Appl. Logic*, to appear.
-  Soskova, A. A. Relativized degree spectra. *J. Logic and Computation* 17, no. 6, 1215-1233 (2007)